



POST-OPERATIVE CARE:

Instructions for Home Care

*It is important to take care of your surgery site.
Please follow these instructions.*

1. Bite on gauze for 30 minutes (one half-hour) to stop any bleeding. If bleeding continues, place another damp gauze and continue to apply pressure.
2. Put ice in a sandwich bag and place ice-bag on cheek or jaw in the area of your extraction. Use the ice for 20 minutes ON and 20 minutes OFF for 2-3 hours.
3. For at least 24 hours DO NOT:
 - smoke
 - drink through a straw
 - drink alcohol, carbonated beverages (soda) or HOT liquids
 - blow your nose
 - spit excessively

Any of these things may cause the blood clot over the surgery site to dislodge from the "socket," causing a "dry socket" and resulting in unnecessary pain. (Should this happen, call this office to see the doctor as soon as possible.)

4. Do maintain a soft diet for the remainder of the day—things like warm soup, ice cream, milk shakes (remember no straws), etc.
5. If your pain, discomfort, or bleeding continues or worsens, call the office for a re-examination.
6. If you were given medications, take them according to the instructions. If you have pain medications at home, you may take them according to the instructions.
7. Tomorrow morning, start rinsing gently with warm salt water (1/4 teaspoon of salt to 1 cup of water) every four (4) hours for two (2) days.
8. If any problems or questions do arise, please feel free to call the office at 228-1085. Our emergency number is available at that number.